

Object-free categories

Stephen A. Fenner

September 30, 2009

Abstract

We define categories where the only primitives are arrows (morphisms) and composition. We recover the objects as identity arrows. These results are known [citation needed].

1 Traditional categories and functors

Definition 1 (Traditional Category). A *category* is a tuple $C = (O, A, \text{dom}, \text{cod}, \circ, \infty)$, where

- O and A are classes (*objects* and *arrows*, respectively),
- $\infty \in A$ (the *undefined arrow* or just *undefined*),
- $\text{dom} : (A - \{\infty\}) \rightarrow O$ and $\text{cod} : (A - \{\infty\}) \rightarrow O$ are functions (*domain* and *codomain*, respectively),
- $\circ : A \times A \rightarrow A$ is a binary (infix) operation on A (*composition*),

and the following axioms are satisfied for all $a \in O$ and $f, g, h \in A$:

1. $\infty \circ f = f \circ \infty = \infty$.
2. $(f \circ g) \circ h = f \circ (g \circ h)$ (composition is associative)
3. $f \circ g \neq \infty$ iff $\text{dom } f = \text{cod } g$.
4. There exists $1_a \in A$ such that $\text{dom } 1_a = \text{cod } 1_a = a$, and $1_a \circ f = f$ if $\text{cod } f = a$ (and $f \neq \infty$), and $g \circ 1_a = g$ if $\text{dom } g = a$ (and $g \neq \infty$). (1_a is an *identity on a*.)

Definition 2 (Traditional Functor). Let $C_1 = (O_1, A_1, \text{dom}, \text{cod}, \circ, \infty)$ and $C_2 = (O_2, A_2, \text{dom}, \text{cod}, \circ, \infty)$ be categories. A *functor* from C_1 to C_2 is a tuple $F = (F_o, F_a)$ such that $F_o : O_1 \rightarrow O_2$ and $F_a : A_1 \rightarrow A_2$ such that the following axioms are satisfied for all $a \in O_1$ and $f, g \in A_1$:

1. $F_a(f) = \infty$ iff $f = \infty$.
2. $F_a(f \circ g) = F_a(f) \circ F_a(g)$.
3. $F_a(1_a) = 1_{F_o(a)}$.

Proposition 1. *If $f \in A_1 - \{\infty\}$, then $F_o(\text{dom } f) = \text{dom } F_a(f)$ and $F_o(\text{cod } f) = \text{cod } F_a(f)$.*

Proof. Let $d = \text{dom } f$ and let $c = \text{cod } f$. We have

$$\infty \neq F_a(f) = F_a(f \circ 1_d) = F_a(f) \circ F_a(1_d) = F_a(f) \circ 1_{F_o(d)}.$$

Thus $\text{dom } F_a(f) = \text{cod } 1_{F_o(d)} = F_o(d) = F_o(\text{dom } f)$. Similarly,

$$\infty \neq F_a(f) = F_a(1_c \circ f) = F_a(1_c) \circ F_a(f) = 1_{F_o(c)} \circ F_a(f).$$

Thus $\text{cod } F_a(f) = \text{dom } 1_{F_o(c)} = F_o(c) = F_o(\text{cod } f)$. □

2 Object-free categories

We will define an *object-free category* to be a tuple $C = (A, \circ, \infty)$, where A is a class (the *arrows*), $\infty \in A$ (*undefined*), and $\circ : A \times A \rightarrow A$ is a binary operator (*composition*) on A . We list the axioms of an object-free category together with some definitions.

In the axioms, f, g, h, \dots are arbitrary elements of A . We say “ f is defined” to mean $f \neq \infty$.

Axiom 1. $f \circ \infty = \infty \circ f = \infty$.

Axiom 2. $(f \circ g) \circ h = f \circ (g \circ h)$.

Axiom 3. *If $f \circ g$ and $g \circ h$ are defined, then so is $f \circ g \circ h$.*

Axiom 4. *If $f_1 \circ g_1$, $f_1 \circ g_2$, and $f_2 \circ g_1$ are all defined, then so is $f_2 \circ g_2$.*

Definition 3. An arrow $i \in A$ is an *identity* (or *object*) if $i \neq \infty$ and for every $f, g \in A$:

- $i \circ f = f$ if $i \circ f$ is defined, and

- $g \circ i = g$ if $g \circ i$ is defined.

Axiom 5. *If f is defined, then there are identities d and c such that $f \circ d$ and $c \circ f$ are both defined. (And both compositions equal f , of course.)*

We now show how to recover the traditional definition from the object-free definition of a category, identifying objects with their identity maps.

Lemma 1. *For any identities i_1 and i_2 , we have $i_1 \circ i_2$ is defined iff $i_1 = i_2$.*

Proof. For the forward direction, if $i_1 \circ i_2$ is defined, then $i_1 = i_1 \circ i_2 = i_2$.

Proving the reverse direction amounts to proving that $i \circ i$ is defined for all identities i . By Axiom 5 with i substituted for f , there is an identity i' such that $i \circ i'$ is defined. But then by the forward direction of the current proposition, $i = i'$, and so $i \circ i$ is defined (and equals i). \square

Proposition 2. *If $f \neq \infty$, then the identities d and c of Axiom 5 are uniquely determined by f .*

Proof. Suppose d_1 and d_2 are identities such that $f \circ d_1$ and $f \circ d_2$ are both defined. By Lemma 1, $d_1 \circ d_1$ is defined. Then by Axiom 4, $d_1 \circ d_2$ is defined, whence $d_1 = d_2$ by Lemma 1 again.

A similar proof shows that if c_1 and c_2 are identities and $c_1 \circ f$ and $c_2 \circ f$ are both defined, then $c_1 = c_2$. \square

Definition 4. *We define $O \subseteq A$ to be the class of all identities (objects). We define the map $\text{dom} : (A - \{\infty\}) \rightarrow O$ to take f to the unique d of Axiom 5. We define the map $\text{cod} : (A - \{\infty\}) \rightarrow O$ to take f to the unique c of Axiom 5. We may also write $f : d \rightarrow c$.*

Theorem 1. *For all $f, g \in A - \{\infty\}$, $f \circ g$ is defined iff $\text{dom } f = \text{cod } g$.*

Proof. For the forward direction, let $d = \text{dom } f$. Then

$$\infty \neq f \circ g = (f \circ d) \circ g = f \circ (d \circ g),$$

which means that $d \circ g$ must be defined (by Axiom 1). Thus $d = \text{cod } g$.

For the reverse direction, suppose $\text{dom } f = \text{cod } g = i$. Then $f \circ i$ and $i \circ g$ are both defined. Hence by Axiom 3, $f \circ i \circ g$ is defined. But $f \circ i \circ g = f \circ g$. \square

Proposition 3. *For any identity i , $\text{dom } i = \text{cod } i = i$.*

Proof. Immediate from the fact that $i \circ i = i$. \square

3 Object-free functors

Definition 5. Let $C_1 = (A_1, \circ, \infty)$ and $C_2 = (A_2, \circ, \infty)$ be object-free categories. An *object-free functor* from C_1 to C_2 is a map $F : A_1 \rightarrow A_2$ such that for all $f, g \in A_1$,

1. $F(f) = \infty$ iff $f = \infty$,
2. $F(f \circ g) = F(f) \circ F(g)$, and
3. if f is an identity, then so is $F(f)$.

We can define $F_a := F$ and F_o to be F restricted to O_1 , noticing that $F(a) \in O_2$ for all $a \in O_1$. (O_1 and O_2 are the classes of objects in A_1 and A_2 , respectively.)

Proposition 4. For any $f \in A_1 - \{\infty\}$, $F(\text{dom } f) = \text{dom } F(f)$ and $F(\text{cod } f) = \text{cod } F(f)$.

Proof. Let $d = \text{dom } f$ and $c = \text{cod } f$. Then $F(d)$ and $F(c)$ are identities, and

$$\begin{aligned}\infty \neq F(f) &= F(f \circ d) = F(f) \circ F(d), \\ \infty \neq F(f) &= F(c \circ f) = F(c) \circ F(f).\end{aligned}$$

Thus $F(d) = \text{dom } F(f)$ and $F(c) = \text{cod } F(f)$. □